



AAP Immunizations Why Immunize?

Disease	What it Does	Year Developed &/or licensed	Why Immunize
Polio	Causes acute paralysis that can lead to permanent physical disability and even death. Visit this page to read a woman recount her struggle with polio at age 3.	1955	Before polio vaccination was available, 13,000 to 20,000 cases were reported each year in the U.S. None were reported in 2010.
Measles	Rash that can cause complications such as pneumonia, diarrhea or ear infections in 9% of those infected. Some develop encephalitis, which results in brain damage. Measles can be fatal, visit this page to read how it claimed the life of an unvaccinated boy in Great Britain.	1963	Between 1958-1962, an average of 503,282 measles cases were reported with 432 deaths. Measles is still one of the most infectious diseases in the world, and is frequently imported into the US. If vaccinations were stopped, 2.7 million measles deaths could be expected worldwide.
Haemophilus Influenzae Type b (Hib) Meningitis	Most common cause of bacterial meningitis in the U.S. before the vaccine. Led to deafness, seizures or mental retardation in those who survived the disease. Click here to learn how Hib devastated the life of one young girl.	1985	Before the vaccination, Hib was the leading cause of childhood bacterial meningitis and postnatal mental retardation. 34 cases of Hib disease were reported among children less than 5 years in 2005. Hib meningitis killed 600 children a year, and infected 20,000. If we were to stop immunizing, we would likely return to the pre-vaccine numbers of infections and deaths.
Pertussis (Whooping Cough)	Can lead to pneumonia, seizures, brain disease and death in infants. Results in prolonged coughing that lasts for many weeks, causing dehydration and vomiting. Listen to Dr. Sarah Long discuss Pertussis.	1926	In 1922, 107,473 cases were reported; 5,099 deaths. Before immunization, up to 260,000 cases were reported in the U.S. each year, with up to 9,000 deaths. Pertussis still occurs worldwide. In 2002, 9,771 cases were reported.
Rubella (German Measles)	Usually mild in children and adults, up to 90% of infants born to infected mothers will develop congenital rubella syndrome (CRS), resulting in heart defects, cataracts, mental retardation and deafness. Read a case study on CRS here.	1969	Before the 1965 vaccination was used routinely in the U.S., rubella resulted in an estimated 20,000 infants born with CRS, 2,100 neonatal deaths and 11,250 miscarriages in a two-year time span. Rubella was declared no longer endemic in US in 2005
Varicella (Chickenpox)	Always present in the community and highly contagious. It can be severe in some and lead to dehydration, pneumonia, and shingles. Children miss a week or more of school on average when sick with chickenpox. Read about 3 unvaccinated children who died from chickenpox.	1995	Chickenpox was responsible for an estimated 4 million cases, 11,000 hospitalizations and 100 deaths each year before the licensing of the chickenpox vaccine. In 1995, 120,624 cases were reported. 22,841 cases were reported in 2002 with 9 deaths.



AAP Immunizations Why Immunize?

<p>Hepatitis B</p>	<p>Infants and children who become infected with Hepatitis B are at the highest risk of developing life-long infection, which often leads to death from liver disease and liver cancer. Listen to Dr. Meg Fisher discuss the importance of HepB vaccine.</p>	<p>1982</p>	<p>In 1982, 22,177 cases were reported. Approximately 25% of children who become infected with life-long hepatitis are expected to die of a related disease as adults. In addition to the 12,000 infants infected by their mother during birth, approximately 33,000 children under the age of 10 were infected before the vaccination. In 2002, 7,996 cases were reported.</p>
<p>Diphtheria</p>	<p>A serious disease caused by poison produced from the bacteria. It frequently causes heart and nerve problems. Diphtheria disease can also be fatal, this story tells of one child's death.</p>	<p>1923</p>	<p>In 1920, 147,991 diphtheria cases were reported and 13,170 deaths. The death rate before vaccinations was up to 20% in the young and elderly. Although Diphtheria is primarily in other countries, international travels make it easy to contract. In 1921, a diphtheria outbreak caused 12,230 deaths in the U.S. 1 case was reported in 2002.</p>
<p>Tetanus (Lock Jaw)</p>	<p>A severe, often fatal disease. Leads to stiffness and spasms of the muscles. Can cause the throat to close, and spasms can cause fractures. To learn more about the risks to those who remain unvaccinated, click here.</p>	<p>1927</p>	<p>Between 1922-1926, an average of 1,314 cases of tetanus were reported per year. Approximately 30% of reported cases of tetanus end in death. Tetanus kills 300,000 newborns and 30,000 birth mothers worldwide, from lack of immunization. Tetanus is not contagious, and can only be prevented by immunization. People of all ages can be infected. From 2001-2008, there were roughly 30 cases reported per year, in the US. 2002, 25 cases were reported in US.</p>
<p>Mumps</p>	<p>Once a major cause of deafness in children, occurring in approximately 1 of every 20,000 cases reported. Can cause swelling of the brain, nerves and spinal cord that can lead to paralysis, seizures and fluid in the brain. This story of a camp counselor who spread mumps to 30 others, shows the importance of everyone being vaccinated.</p>	<p>1967</p>	<p>Before the vaccination was developed in 1967, an estimated 212,000 cases occurred in the U.S. annually. In 1986 and 1987, there was a resurgence of mumps with 12,848 cases reported. Since 1989, the incidence has declined, with a total of 323 cases last year. From 2009-2010, another outbreak of 3,502 cases occurred in the US.</p>
<p>Smallpox</p>	<p>Smallpox is a serious, contagious, and sometimes fatal infectious disease. There is no specific treatment for smallpox disease, and the only prevention is vaccination. The <i>pox</i> part of <i>smallpox</i> is derived from the Latin word for "spotted" and refers to the raised bumps that appear on the face and body of an infected person.</p>	<p>1796 (not widely used before 1900)</p>	<p>In 1900, 21,064 smallpox cases were reported; 894 deaths. Smallpox was eradicated in 1977</p>



AAP Immunizations Why Immunize?

<p>Influenza</p>	<p>The flu is a contagious respiratory illness caused by influenza viruses. It can cause mild to severe illness, and at times can lead to death. Some people, such as older people, young children, and people with certain health conditions (such as asthma, diabetes, or heart disease), are at high risk for serious flu complications. Read this family story on how a young boy was hospitalized due to the flu.</p>	<p>1945 (not routinely used until 2004)</p>	<p>Every year in the United States, on average 5% to 20% of the population gets the flu; more than 200,000 people are hospitalized from flu-related complications; and about 36,000 people die from flu-related causes.</p>
<p>Pneumococcal</p>	<p>Symptoms include pneumococcal pneumonia (high fever, cough, and shortness of breath), bacteremia (fever and feeling generally poorly), and meningitis (fever, headache, thinking slowly or not clearly). Read this family story about a baby boy who contracted pneumococcal meningitis.</p>	<p>1977 (polysaccharide) 2000 (conjugate – childhood vaccine)</p>	<p>In 1998-1999, rate of invasive pneumococcal disease in children was 188 per 100,000 population. Disease rates have declined 70-80% among children younger than 2 years of age, compared to 1998-1999.</p>
<p>Hepatitis A</p>	<p>Hepatitis A is a serious liver disease. Hepatitis A can cause flu-like illness, jaundice, severe stomach pains and diarrhea (children). Some who contract Hepatitis A will die each year. To learn how debilitating this virus can be, read this story.</p>	<p>1995</p>	<p>Hepatitis A rates have been declining since 1995, and since 1998 have been at historically low levels. A total of 1,987 cases was reported in 2009. The wider use of vaccine is probably contributing to this marked decrease in hepatitis A rates in the United States.</p>
<p>Human Papillomavirus (HPV)</p>	<p>HPV is the most common sexually transmitted virus in the United States. HPV can cause cervical cancer in women, and is associated with several less common cancers, in both men and women. This is the story of young woman who lost her fertility to cervical cancer.</p>	<p>2006 (Gardasil, now for males and females) 2009 (Cervarix for females only)</p>	<p>An estimated 20 million persons are currently infected, and an estimated 6.2 million new HPV infections occur annually. HPV infection is common among adolescents and young adults. Prevalence among adolescent girls is as high as 64%. Up to 75% of new infections occur among persons 15–24 years of age. Modeling estimates suggest that more than 80% of sexually active women will have been infected by age 50.</p>
<p>Rotavirus</p>	<p>Rotavirus is a virus that causes diarrhea (sometimes severe), mostly in babies and young children. It is often accompanied by vomiting and fever, and can lead to dehydration. To read a story of rotavirus infection, click here.</p>	<p>2006</p>	<p>In the prevaccine era rotavirus infection was responsible for more than 400,000 physician visits, more than 200,000 emergency department (ED) visits, 55,000 to 70,000 hospitalizations each year, and 20 to 60 deaths. Rotavirus activity in the United States decreased significantly after introduction of rotavirus vaccine in 2006.</p>



AAP Immunizations Why Immunize?

<p>Meningococcal</p>	<p>Meningococcal bacteria is another cause of Meningitis, a serious infection of the covering of the brain. It can also cause blood infection. 10-15% of Meningitis cases cause death, of survivors 11-19% will lose their arms and/or legs, become deaf, mentally retarded, or suffer seizures or strokes. Read this family story to learn how quickly Meningitis took the life of a teenage boy.</p>	<p>1974 (MPSV4) 2010 (Menveo)</p>	<p>Prior to 2000, an estimated 1,400 to 2,800 cases of meningococcal disease occurred each year in the United States, a rate of 0.5 to 1.1 per 100,000 population. Incidence has decreased since 2000, and incidence of serogroups C and Y, which represent the majority of cases of vaccine-preventable meningococcal disease, are at historic lows.</p>
--------------------------------------	--	---------------------------------------	---

Source: Centers for Disease Control and Prevention (www.cdc.gov)

References:

- Centers for Disease Control and Prevention. Achievements in Public Health, 1900-1999. Impact of Vaccines Universally Recommended for Children – United States, 1990-1998. *Morbidity and Mortality Weekly Report*. April 2, 1999;48:243-248. Available at: <http://www.cdc.gov/mmwr/PDF/wk/mm4812.pdf>
- Centers for Disease Control and Prevention. Summary of Notifiable Diseases – United States, 2002. *Morbidity and Mortality Weekly Report*. April 30, 2004;51:1-84. Available at: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5153a1.htm>
- Centers for Disease Control and Prevention. What Would Happen If We Stopped Vaccinations. Available at: <http://www.cdc.gov/vaccines/vac-gen/whatifstop.htm>
- Centers for Disease Control and Prevention. Epidemiology and Prevention of Vaccine-Preventable Diseases, The Pink Book, 8th Ed. January 2005. Available at: <http://www.cdc.gov/vaccines/pubs/pinkbook/default.htm>
- Centers for Disease Control and Prevention. Epidemiology and Prevention of Vaccine-Preventable Diseases, The Pink Book, 12th Ed. February 2012. Available at: <http://www.cdc.gov/vaccines/pubs/pinkbook/index.html>

